



Application of Blue-green Alga *Spirulina* for removing Caesium ions from polluted water

M. Kurashvili*, G. Adamia, T. Varazi, G. Khatisashvili, G. Gigolashvili, M. Pruidze, L. Chokheli, S. Japharashvili

Durmishidze Institute of Biochemistry and Biotechnology of Agricultural University of Georgia, 240, David Agmashenebeli, Alley, Tbilisi, 0159, Georgia;

Received: 23 October 2018; accepted: 12 December 2018

ABSTRACT

The presented work concerns studying a possibility of using blue-green alga (cyanobacteria) *Spirulina* (*Spirulina platensis*) for removal of Caesium ions from polluted waters. For this aim, tolerance of alga to Cs^+ ions, in particular, influence of different concentrations of heavy metal on ultrastructure of *Spirulina* cells, as well as on biomass production by *Spirulina* and chlorophyll accumulation in alga cells have been studied. According to obtained data, the highest concentration of Cs^+ (100 ppm) blocked biomass formation only by 25%, and the content of chlorophyll decreased by 20%. This concentration of heavy metal caused only an increase of hydrocarbons-containing space between lamella in *Spirulina* cells, which can be a result of accumulation of Cs^+ ions in alga cells. Thus, 100 ppm concentration of Cs^+ ions has been chosen for testing possibility of using *Spirulina* to clean artificially polluted water in the model large-scale experiment. The obtained data have shown that *Spirulina* has capability to assimilate up to 90% of Cs^+ ions from polluted water during 15 days. These results indicate that *Spirulina* is potentially useful for cleaning waters polluted with nonradioactive Cs^+ ions as well as with $^{137}\text{Cs}^+$ ions.

Keywords: Phytoremediation, *Spirulina platensis*, Water pollution, Heavy metals, Caesium ions, Radioactive and Stable Cesium.

*Corresponding author: Maritca Kurashvili, E-mail address: m.kurashvili@agrundi.edu.ge

1. Introduction

Nowadays, water pollution by different chemicals especially by hazardous heavy metals and radionuclides is one of the most important ecological problem. Among them ^{137}Cs is very dangerous. Nonradioactive Caesium ions that occur in nature and are released into environment through mining and milling of ores, are only mildly toxic [1]. Both radioactive and stable cesium, when they occur in human or animal body, act the same way [2].

Presented work devotes to elaboration of method for cleaning water contaminated with Cs^+ ions by using blue-green alga *Spirulina* (*Spirulina platensis*). *Spirulina* (*Spirulina platensis*) should have prospects for phytoremediation of waters

polluted by different toxic compounds. *Spirulina* has unique chemical content and biological properties. This blue-green alga is characterized by reproduction and fast biomass formation in extreme conditions. *Spirulina* cells have rich content of proteins and peptides, and these compounds should be chelating ions of heavy metals. During the aging, the vacuoles of *Spirulina* cells inflate with air bubbles and as a result, *Spirulina* colonies float onto the water surface. As a result, *Spirulina* biomass will be easily (mechanically) separated from cleared water body after remediation. These factors have attracted our interest in *Spirulina* as to potential phytoremediator agent. It is known the information about using of algae for cleaning the environment polluted with such toxicants as heavy metals [3-

8], radioactive elements [9], fluoride ions [10] and others. There is only limited data concerning using *Spirulina* as a phytoremediator agent in case of water pollution with Cs^+ ions, but for this aim, *Spirulina* seems to have prospects, because there is a lot of information about heavy metals uptake processes by *Spirulina* [3-5].

In presented work, potential of *Spirulina* to uptake of Cs^+ ions from water solutions has been evaluated. For this aim, tolerance of alga to Cs^+ ions, in particular, the influence of different concentrations of heavy metal on ultrastructure of *Spirulina* cells, as well as on biomass formation and chlorophyll accumulation by *Spirulina* have been studied. According to the obtained data, the phytoremediation technology for cleaning of water polluted with Caesium based on application of *Spirulina* have been elaborated and tested via model experiments.

2. Materials and methods

2.1 Materials

The biomass of *Spirulina platensis* obtained via cultivation in standard Zarrouk's medium (pH – 8.7; content in g/L: NaHCO_3 – 16.8, K_2HPO_4 – 0.5, NaNO_3 – 2.5, K_2SO_4 – 1.0, NaCl – 1.0, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ – 0.2, $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ – 0.04, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ – 0.01, EDTA – 0.08; and microelements kit A5 – 1 mL) were used in the experiments. Incubation was carried out with permanent air barbotage (rate of air flow 2 L/min), at temperature 25°C, and under following illumination conditions: a photoperiod of lighting 16L/8D (16 hours of light: 8 hours of dark), a total photosynthetic photon flux density (PPFD) of $\approx 15 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$.

2.2 Methods

For measurement of fresh biomass productivity and chlorophyll formation by *Spirulina*, the following method was elaborated: the incubation medium was centrifuged at 1000 g during 20 min and obtained pellet was weighted. The obtained fresh biomass was treated by acetone and the content of chlorophyll was determined spectrophotometrically at 652 nm according to standard method [11].

The biomass of *Spirulina* in incubation medium have been measured spectrophotometrically at 750 nm [12].

The content of Cs^+ ions in the samples was determined by the method of atomic absorption

(flame emission) analysis [13]. Conditions for analysis are the following: Instrument – PerkinElmer HGA900 Graphite Furnace; wavelength – 852.1 nm; slit – 0.2/0.4 nm; flame – air-acetylene; stock standard solution – CESIUM 1000 mg/L; light source – EDL; interference – ionization controlled by addition of 0.1% KCl.

To study the influence of Cs^+ ions on cell ultrastructure, biomass of *Spirulina* was fixed in 2.5% glutaraldehyde and then in 1% OsO_4 . After dehydration in ethanol of increasing concentrations, the samples were embedded in Epon–Araldite resin (1.5: 1.0) and poured into gelatin capsules [14]. Thin serial sections (50–60 nm) were made using an LKB III ultra-microtome, stained with 2% uranyl acetate, and analyzed by electron microscope Tesla BS 500 (Czech Republic) (specifics, transmission; resolution, 0.8 nm).

2.3 Experimental design

For estimation of influence of different concentrations of Cs^+ on growing parameters of *Spirulina*, alga was cultivated in standard Zarrouk's media (volume 50 mL) containing different concentrations of Cs^+ (1, 10 or 100 ppm). The initial biomass of *Spirulina* was 3-4 g/L. The incubation was carried out in flasks during 72 h (120 h for ultrastructural investigations), at temperature 25°C, without air barbotage, a photoperiod of lighting 16L/8D, PPFD $\approx 15 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$. After incubation the biomass productivity by *Spirulina* and chlorophyll content in cells by above mentioned methods were determined. The ultrastructure analysis was carried out in same samples.

The model large-scale experiment was carried out in following conditions: *Spirulina* was preliminary growing in Zarrouk's medium (volume 20 L) and after 7 days 20 L of Cs^+ containing solution was added. The volume of incubation medium in the beginning of incubation of *Spirulina* with Cs^+ was 40 L. At that moment, the biomass of *Spirulina* was 3.5 g/L and Cs^+ concentration was 100 ppm. The incubation was carried out in a glass container with sizes 60 x 21 x 40 (in cm, length x width x height), with permanent air barbotage (rate of air flow 2 L/min), at temperature 25°C, under following illumination conditions: a 24-h lighting (24L/0D), PPFD $\approx 15 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$. In the control variant instead of Cs^+ solution 20 L of water was added. The content of Cs^+ ions and biomass of *Spirulina* were determined after 5, 10 and 15 days from addition of heavy metal ions.

3. Results and Discussion

3.1. The influence of Cs^+ on *Spirulina* cells

The influence of different concentrations of Caesium ions (1, 10 and 100 ppm) on biomass and chlorophyll accumulation by *Spirulina* has been studied. The obtained results are given in Fig. 1.




Fig. 1. The influence of different concentrations of caesium ions on biomass formation and chlorophyll accumulation by *Spirulina*. Incubation conditions are described in part 2.3.

According to the obtained results, low concentrations of heavy metal revealed insignificant effect on *Spirulina* vital processes - reduction of biomass formation (by 5-6%) and chlorophyll content (by 8-9%). The high concentration of Cs^+ (100 ppm) blocked biomass production by 25%, and the content of chlorophyll decreased by 20%.

According to the obtained results, *Spirulina* revealed tolerance to all tested concentrations of Cs^+ ions.

3.2. Study of penetration and localization of Cs^+ in *Spirulina*

On the next stage, the influence of Cs^+ ions on ultrastructure of *Spirulina* cells using electron microscopic method has been studied (Fig. 2 A-D).

As Fig. 2 shows, the influence of the highest concentration of Cs^+ ions (100 ppm) caused only an increase of hydrocarbons-containing space between lamella in *Spirulina* cells, which can be a result of accumulation of heavy metal ions. Thus, 100 ppm concentration of Cs^+ ions has been chosen for the model experiments.

3.3. The results of model experiment

The model experiments for elaboration new approach of phytoremediation technology for cleaning water polluted by Cs^+ ions have been carried out. For this aim, *Spirulina* cultivated in water artificially polluted with 100 ppm Cs^+ ions. The obtained data are given in Fig. 3.

As it seen from Fig. 3, *Spirulina* has capability to assimilate up to 90% of heavy metal ions from polluted water during 15 days. At the same time biomass of alga decreases by 25%. These results indicate that *Spirulina* is potentially usable for cleaning of waters polluted with nonradioactive Cs^+ ions as well as with $^{137}\text{Cs}^+$ ions.




Fig. 2. Electron micrographs of cells of *Spirulina*, cultivated on Zarrouk's medium without Cs^+ ions (A and B), and with 100 pm of Cs^+ ions. Incubation conditions are described in part 2.3.




Fig. 3. The results of model experiment for cleaning Cs⁺ containing water by using Spirulina. The conditions of experiment are described in part 2.3.

5. Conclusions

Thus, the work carried out has shown that Spirulina cells have an ability to effectively remove Cs⁺ ions from artificially polluted water. These results can become a basis for development of a new phytoremediation technology for cleaning waters polluted with radioactive Caesium ions (¹³⁷Cs⁺) based on application of Spirulina.

6. Acknowledgment

This work was supported by Shota Rustaveli National Science Foundation (SRNSF), with cofounding by Georgian Agricultural University - Applied Research Grant # 216944 and SRNSF - CNR # 04/51.

References

- [1] C. Pinsky, R. Bose, J.R. Taylor, J. McKee, C. Lapointe, J. Birchall, Cesium in mammals:

Acute toxicity, organ changes and tissue accumulation, ". J. Environ. Sci. Health. Part A, 16 (1981) 549–567.

- [2] J. Rundo, A Survey of the Metabolism of Caesium in Man, British J. Radiol. 37 (1964) 108–114.
- [3] A. Ahmad, R. Ghufuran, Z.A. Wahid, Cd, As, Cu, and Zn Transfer through Dry to Rehydrated Biomass of Spirulina Platensis from Wastewater, Polish J. Environ. Stud. 19 (2010) 887–893.
- [4] H. Chen, S. Pan, Bioremediation potential of spirulina: toxicity and biosorption studies of lead, J. Zhejiang Univ. Sci. B. 6 (2005) 171–174.
- [5] O. Murali, S. Mehar. Bioremediation of heavy metals using Spirulina. Int. J. Geology, Earth and Environ. Sci. 4 (2014) 244–249.
- [6] K. Suresh Kumar, H.-U. Dahms, E.-J. Won, J.-S. Lee, K.-H. Shin, Microalgae – A promising tool for heavy metal remediation, Ecotoxicol. Environ. Saf. 113 (2015) 329–352.

- [7] A.H. Mahvi, Cadmium Biosorption from Wastewater by Ulmus Leaves and Their Ash, *Europ. J. Scient. Res.* 23 (2008) 197-203.
- [8] A. Maleki, A.H. Mahvi, M.A. Zazouli, H. Izanloo, A.H. Barati, Aqueous Cadmium Removal by Adsorption on Barley Hull and Barley Hull Ash, *Asian J. Chem.* 23 (2011) 1373-1376.
- [9] S. Fukuda, K. Lwamoto, M. Asumi, A. Yokoyama, T. Nakayama, K. Ishida, I. Inouye, Y. Shraiwa, Global searches for microalgae and aquatic plants that can eliminate radioactive cesium, iodine and strontium from the radio-polluted aquatic environment, *J. Plant Res.* 127 (2014) 79-89.
- [10] M.A. Zazouli, A.H. Mahvi, S. Dobaradaran, M. Barafrashtehpour, Y. Mahdavi, D. Balarak, Adsorption of fluoride from aqueous solution by modified *Azolla filiculoides*, *Fluoride* 47 (2014) 349-358.
- [11] D.I. Arnon, Copper enzymes in isolated chloroplasts, Polyphenoxidase in *Beta Vulgaris*. *Plant Physiol.* 24 (1949) 1-15.
- [12] C. Butterwick, S.I. Heaney, J.F. Talling. A comparison of eight methods for estimating the biomass and growth of planktonic algae, *British Phycol.J.*, 17, (1982) 69-79.
- [13] L.V. Guba, I.I. Dovgvy, M.A. Rizhkova, Method of measuring of cesium by flame emission photometry method. *Scientific Notes of Taurida V.Vernadsky National University. Series: Biol. Chem.* 25 (2012) 284-288.
- [14] O. Buadze, T. Sadunishvili, G. Kvesitadze, The effect of 1,2-benzanthracene and 3,4-benzpyrene on the ultrastructure of maize cells, *Int. Biodeter. Biodegr.* 44 (1998) 49–54.